arXiv:math/0607206v4 [math.PR] 15 May 2008

Dualities for Multi-State Probabilistic Cellular Automata

F. Javier López[1], **Gerardo Sanz**[1] and **Marcelo Sobottka**[2]

[1]Departamento de Métodos Estadísticos and BIFI, Facultad de Ciencias, Universidad de Zaragoza, C/ Pedro Cerbuna 12, 50009 Zaragoza, Spain.
[2]Departamento de Ingeniería Matemática, Facultad de Ciencias Físicas y Matemáticas, Universidad de Concepción, Casilla 160-C, Correo 3, Concepción, Chile.

E-mail: `javier.lopez@unizar.es`, `gerardo.sanz@unizar.es`, `msobottka@udec.cl`

Abstract.

In this paper a new form of duality for probabilistic cellular automata (PCA) is introduced. Using this duality, an ergodicity result for processes having a dual is proved. Also, conditions on the probabilities defining the evolution of the processes for the existence of a dual are provided. The results are applied to wide classes of PCA which include multi-opinion voter models, competition models and the Domany-Kinzel model.

Keywords: cellular automata; stochastic particle dynamics (theory); interacting agent models; stochastic processes.

1. Introduction

Probabilistic Cellular Automata (PCA) are discrete-time stochastic processes with state space $\mathbf{X} := W^{\mathbb{Z}^d}$, where $W = \{1, \ldots, M\}$ and $d \geq 1$, with finite range interactions on the integer lattice $\mathbb{Z}^d$ (this is interpreted as at each point $z \in Z^d$ there is a particle which take values in the set W; the points $x \in \mathbf{X}$ are called configurations). More precisely, suppose $N \subset \mathbb{Z}^d$ is a finite subset and $f : W^N \times W \to [0, 1]$ is a transition function, then a PCA is a discrete-time homogenous Markov process $\eta_t = \{\eta_t(z) \in W : \ z \in \mathbb{Z}^d\}$ whose evolution is given, for all $s \in \mathbb{N}$, $z \in \mathbb{Z}^d$, $w \in W$, and $x \in \mathbf{X}$, by

$$\mathbb{P}\left\{\eta_{s+1}(z) = w | \ \eta_s = x\right\} = f((x(z+n))_{n\in N}, w). \tag{1}$$

In other words, η_t is an interacting particle system on $\mathbf{X}$ in discrete time such that the transitions at different sites are conditionally independent, given the current state of the system. For definitions and main results on PCA, the reader can consult [1] and [2].

The set $\mathbf{X} = W^{\mathbb{Z}^d}$ is endowed with the product topology. Let $\mathcal{P}(\mathbf{X})$ be the set of probability measures on $\mathbf{X}$ with the weak* convergence topology. Note that $\mathcal{P}(\mathbf{X})$ is compact with respect to this topology.

Let η_t and ξ_t be two stochastic processes (which can evolve either in continuous or discrete time) with state spaces $\mathbf{X}$ and $\mathbf{Y}$ respectively, and suppose $H : \mathbf{X} \times \mathbf{Y} \to \mathbb{R}$ is a bounded measurable function, then η_t and ξ_t are dual to one another with respect to H if for all $x \in \mathbf{X}$ and $y \in \mathbf{Y}$ (Definition II.3.1 of [3])

$$\mathbb{E}_{\eta_0=x}\left[H(\eta_s, y)\right] = \mathbb{E}_{\xi_0=y}\left[H(x, \xi_s)\right], \tag{2}$$

where $\mathbb{E}_{\gamma_0=\gamma}[\cdot]$ denotes the expectation when the process γ_t starts with configuration γ.

If η_t and ξ_t are both discrete-time Markov chains with transition matrices P and Q respectively, then the previous expression for duality can be written as

$$P^s H = H(Q^s)^T, \tag{3}$$

where the superscript ^{T} stands for the transpose. Furthermore, due to the Markovian property, if the duality equation holds for $s = 1$ then it holds for all $s \in \mathbb{N}$.

Duality allows obtaining relevant information about the evolution of η_t, which usually has uncountable state space, by studying the evolution of ξ_t which in general is chosen having countable state space. In fact, duality has been widely used in the study of spin systems (continuous-time interacting particle systems with $W = \{0, 1\}$), where $\mathbf{Y}$ is the set of all finite subsets of $\mathbb{Z}^d$ (see [3], [4], [5], and [6]). For the case where η_t is a continuous-time interacting particle system with $|W| \geq 2$, López and Sanz [7] studied an extended version of (2), which is given by the equation:

$$\mathbb{E}_{\eta_0=x}\left[H(\eta_s, y)\right] = \mathbb{E}_{\xi_0=y}\left[H(x, \xi_s)e^{-\int_0^s V(\xi_u)du}\right], \tag{4}$$

for all $x \in \mathbf{X}$ and $y \in \mathbf{Y}$, and for some function $V : \mathbf{Y} \to [0, \infty)$. In particular, this duality equation was used in [7] to obtain results about the long time behaviour of the continuous-time multi-opinion noisy voter model and the continuous-time 3-opinion noisy biased voter model.

For the discrete-time case, Katori et al. [8] develop a theory of duality for two state PCA based on the equation

$$\mathbb{E}\left[\mathfrak{c}^{|\xi_s^B \cap A|}\right] = \mathbb{E}\left[\mathfrak{c}^{|\eta_s^A \cap B|}\right], \tag{5}$$

where $\mathfrak{c}$ is a parameter depending on the processes η_t and ξ_t; $A, B \subseteq \mathbb{Z}$, with at least one of them being finite; η_s^A and ξ_s^B represent the set of particles with value 1 at time s starting from configurations where only the particles in A and B, respectively, have the value 1; and $|\cdot|$ stands for the cardinality of a set (this equation was first considered for interacting particle systems by Sudbury and Lloyd in [9]). In [8], this relationship is used to study duality for some cases of the Domany-Kinzel model, which is the PCA with state space $\{0, 1\}^{\mathbb{Z}}$, and evolution defined by the parameters $a_0, a_1, a_2 \in [0, 1]$ as follows:

$$\mathbb{P}\left\{\eta_{s+1}(z) = 1 |\ \eta_s(z-1) = w_{-1}, \eta_s(z) = w_0, \eta_s(z+1) = w_1\right\} = a_{w_{-1}+w_1} \tag{6}$$

When $a_0 = 0$ and $a_2 \leq a_1$, Katori et al. [8] have presented a dual for η_t. Such dual is a model in which each stage is updated by a a_1-thinning of all sites, followed by an application of the Domany-Kinzel model with parameters $a_0' = 0$, $a_1' = 1$, and $a_2' = a_2/a_1$. When $a_1 > a_2$, they also studied the limit behaviour of the process. The technique used in [8] was to locate the process η_t in a finite subset of $\mathbb{Z}$ to represent it by a transition matrix P, and so to find H and Q which satisfy the duality equation (3). The technique used by Katori et al. depends strongly on the fact that $a_0 = 0$ (see also chapter 5 of [4] for results on this model, with $a_0 = 0$, using percolation theory). In [10] and [11], those results are extended to a larger class of one dimensional two state PCA.

For multi-state PCA, an equation analogous to (5) is introduced in [12] to define self-dual processes (that is, processes η_t that satisfy the equation with $\xi_t = \eta_t$) and give conditions for self-duality.

Our objective in this paper is to give a general theory of duality for multi-state PCA. The duality equation we define (see (7) below) is given for general duality functions H and is not restricted to self-duality. Moreover, the introduction of an auxiliary function d in the equation allows the inclusion of wider classes of processes for each dual function H. We will show the usefulness of our concept by obtaining an ergodicity result (Theorem 2.1) for processes satisfying (7). Equation (7) and Theorem 2.1 are stated for general H and we show their applicability with two particular cases of H which cover two wide classes of PCA, providing conditions on their transition probabilities such that they satisfy (7) and, in that case, providing conditions for their ergodicity (that is, the existence of an unique equilibrium measure and the convergence of the process to that measure). We will also apply our results to some examples.

The duality equation we propose is

Definition 1.1. *Given two discrete-time Markov processes, η_t with state space $\mathbf{X}$ and ξ_t with state space $\mathbf{Y}$, and $H : \mathbf{X} \times \mathbf{Y} \to \mathbb{R}$ and $d : \mathbf{Y} \to [0, \infty)$ bounded measurable functions, we say η_t and ξ_t are dual to one another with respect to (H, d) if*

$$\mathbb{E}_{\eta_0 = x}\left[H(\eta_1, y)\right] = d(y)\mathbb{E}_{\xi_0 = y}\left[H(x, \xi_1)\right]. \tag{7}$$

Notice that the duality equation with respect to (H, d) is only written for the one-step evolution of the processes η_t and ξ_t as they are both Markov processes, but an induction argument yields the duality equation for the s step evolution of the process:

$$E_{\eta_0 = x}[H(\eta_s, y)] = d(y)E_{\xi_0 = y}[d(\xi_1)\cdots d(\xi_{s-1})H(x, \xi_s)]. \tag{8}$$

When η_t and ξ_t are Markov chains with transition matrix P and Q, respectively, equation (7) can be written as

$$PH = H(DQ)^T, \tag{9}$$

where D is the diagonal matrix with $D_{yy} = d(y)$. In this case, the general expression for time $s \geq 1$ is $P^s H = H\left((DQ)^s\right)^T$.

The advantage of considering the function d on the right side of (7) is that processes that do not have dual with respect to some function H in (2) may have it with respect to (H, d) (see Remarks 3.9 and 3.15 below). When $d \equiv 1$, we have the classical notion of duality (2), which means that the evolution of η_t is easily understood from the evolution of ξ_t. When $d \not\equiv 1$, the relationship between η_t and ξ_t becomes more complicated; nevertheless, in that case, important information about η_t can be obtained from ξ_t.

Although equations (7) and (8) can be seen as a discrete counterpart of equation (4), given for the continuous-time case, there are several differences between the techniques used to exploit the former and the latter. In fact, instead of using directly equation (8) to prove results on ergodicity for the process, we show that there is a transformation $\tilde{\xi}_t$ of the dual process ξ_t such that η_t and $\tilde{\xi}_t$ are dual in the classical sense (with respect to a modified function $\tilde{H}$), see Theorem 2.1(a) below. This approach is useful, for instance, for writing down the explicit form of the invariant measure (11) and for providing alternative conditions for ergodicity (see Remark 2.2) which are used, for instance, in the study of the Domany-Kinzel model (Corollary 3.8).

The organization of the paper is as follows. In §2 we present the basic tools for the duality of stochastic processes and obtain conditions on H and d for the ergodicity of PCA. In §3 we analyze, using the duality theory developed, two general classes of PCA. For each of these classes, we first obtain conditions on the transition probabilities of the processes so that they have a dual and then give conditions for their egodicity. The results are applied to specific models such as the multi-opinion noisy voter model, the Domany-Kinzel model and competition models. For the particular case of the Domany-Kinzel model with parameters $0 \leq a_0 \leq a_1 \leq a_2$ we find a dual process for many cases not covered by [4] or [8]; and we get new results for the ergodicity of this model.

2. Dualities

In this section we consider probabilistic cellular automata η_t for which there exist a pair of functions (H, d) and a dual process ξ_t with respect to (H, d). Sufficient conditions on (H, d) for the ergodicity shall be presented together with the characterization of the corresponding equilibrium measure.

In what follows, we first recall some basic definitions and results about dualities for stochastic processes. Suppose η_t on $\mathbf{X}$ and ξ_t on $\mathbf{Y}$ are two stochastic processes that are dual to one another with respect to a function H. Given a probability measure μ on $\mathbf{X}$, define for any $y \in \mathbf{Y}$

$$\hat{\mu}(y) := \int_{\mathbf{X}} H(x, y) d\mu(x).$$

For any $s \geq 0$, denote by μ_s the distribution obtained from the initial distribution μ when η_t evolves until time s. In our setting, $\mathbf{X} = W^{\mathbb{Z}^d}$ and η_t is a PCA given by (1); then μ_s is defined on the cylinder $\mathbf{w} = [w_{z_1}, \ldots, w_{z_m}] := \{x \in \mathbf{X} : \; x(z_i) = w_{z_i}, 1 \leq i \leq m\}$ by

$$\mu_s(\mathbf{w}) = \sum_{\mathbf{u} \in \mathcal{C}_s} \mu(\mathbf{u}) \mathbb{P}_{\eta_0 \in \mathbf{u}} \{\eta_s \in \mathbf{w}\},$$

where $\mathcal{C}_s$ is the family of all cylinders of $\mathbf{X}$ defined on the coordinates $\{z_i + n_j : \; i = 1, \ldots, m; \; n_j \in N\}$.

Therefore, we have that

$$\hat{\mu}_s(y) = \int_{\mathbf{X}} H(x,y) d\mu_s(x) = \int_{\mathbf{X}} \mathbb{E}_{\eta_0=x}\left[H(\eta_s,y)\right] d\mu(x),$$

and by the duality equation (2) it follows that

$$\begin{aligned} \hat{\mu}_s(y) &= \textstyle\int_{\mathbf{X}} \mathbb{E}_{\eta_0=x}\left[H(\eta_s,y)\right] d\mu(x) \\ &= \textstyle\int_{\mathbf{X}} \mathbb{E}_{\xi_0=y}\left[H(x,\xi_s)\right] d\mu(x) \\ &= \mathbb{E}_{\xi_0=y}\left[\textstyle\int_{\mathbf{X}} H(x,\xi_s) d\mu(x)\right] \\ &= \mathbb{E}_{\xi_0=y}\left[\hat{\mu}(\xi_s)\right]. \end{aligned} \tag{10}$$

Suppose the set of the linear combinations of the functions $\{H(\cdot,y):\ y \in \mathbf{Y}\}$ is dense in $C(\mathbf{X})$, the space of continuous real functions on $\mathbf{X}$. If for any y, $E_{\xi_0=y}[\hat{\mu}(\xi_s)]$ converges to some function of y (not depending on μ) as s goes to ∞, then for any $f \in C(\mathbf{X})$, it follows that $\int_{\mathbf{X}} f d\mu_s$ converges to some functional $\tilde{\nu}(f)$, which means μ_s converges in the weak* topology to some probability measure ν. Furthermore, if the above holds for any initial distribution μ, it implies the ergodicity of η_t: that is, the convergence in distribution of the process to the equilibrium measure ν from any initial measure.

Now we are able to study the case of the duality with respect to (H,d).

Theorem 2.1. *Suppose η_t is a Markov process with state space $\mathbf{X}$ and ξ_t is a Markov chain with state space $\mathbf{Y}$, which are dual to one another with respect to (H,d), where $0 \le d(y) \le 1$ for all $y \in \mathbf{Y}$. Then:*

(a) *There exist a Markov chain $\tilde{\xi}_t$ with state space $\tilde{\mathbf{Y}} = \mathbf{Y} \cup \{\wp\}$, where $\wp$ is a new state, and a bounded measurable function $\tilde{H} : \mathbf{X} \times \tilde{\mathbf{Y}} \to \mathbb{R}$ such that η_t and $\tilde{\xi}_t$ are dual to one another with respect to $\tilde{H}$;*

(b) *denoting by Θ the set of all absorbing states of ξ_t, if*

(i) the linear combinations of $\{H(\cdot,y):\ y \in \mathbf{Y}\}$ is a dense set of $C(\mathbf{X})$;

(ii) $d(y) < 1$ for any $y \notin \Theta$, and $\sup_{y\in\mathbf{Y}:\ d(y)<1} d(y) < 1$;

(iii) $H(\cdot,\theta) \equiv c(\theta)$ for all $\theta \in \Theta$ with $d(\theta) = 1$;

then η_t is ergodic and its unique equilibrium measure is determined for any $y \in \mathbf{Y}$ by

$$\hat{\nu}(y) = \sum_{\substack{\theta \in \Theta, \\ d(\theta)=1}} c(\theta)\mathbb{P}_{\tilde{\xi}_0=y}\left\{\tilde{\xi}_\tau = \theta\right\}, \tag{11}$$

where τ is the hitting time of $\{\theta \in \Theta:\ d(\theta)=1\} \cup \{\wp\}$ for $\tilde{\xi}_t$.

Proof. **(a)** Since $0 \leq d(y) \leq 1$ for all $y \in \mathbf{Y}$, we can define a Markov chain $\tilde{\xi}_t$ with state space $\tilde{\mathbf{Y}} = \mathbf{Y} \cup \{\wp\}$ and transition probabilities given by

$$\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\{\tilde{\xi}_1 = \tilde{y}'\} := \begin{cases} d(\tilde{y})\mathbb{P}_{\xi_0=\tilde{y}}\{\xi_1 = \tilde{y}'\} & , \ if \quad \tilde{y}, \tilde{y}' \in \mathbf{Y}, \\ \\ 1 - d(\tilde{y}) & , \ if \quad \tilde{y} \in \mathbf{Y}, \ \tilde{y}' = \wp \\ \\ 1 & , \ if \quad \tilde{y} = \tilde{y}' = \wp \end{cases} \tag{12}$$

Note that the new state $\wp$ is an absorbing state of the process $\tilde{\xi}_t$.

Define $\tilde{H} : \mathbf{X} \times \tilde{\mathbf{Y}} \to \mathbb{R}$ by

$$\tilde{H}(x, \tilde{y}) := \begin{cases} H(x, \tilde{y}) & , \ if \quad \tilde{y} \in \mathbf{Y}, \\ \\ 0 & , \ if \quad \tilde{y} = \wp \end{cases}$$

It is trivial that (2) holds for $x \in \mathbf{X}$ and $\wp$ since $\tilde{H}(x, \wp) = 0$. On the other hand, if $\tilde{y} \in \mathbf{Y}$, then

$$\begin{aligned} \mathbb{E}_{\eta_0=x}\left[\tilde{H}(\eta_1, \tilde{y})\right] &= \mathbb{E}_{\eta_0=x}\left[H(\eta_1, \tilde{y})\right] \\ &= d(\tilde{y})\mathbb{E}_{\xi_0=\tilde{y}}\left[H(x, \xi_1)\right] \\ &= d(\tilde{y}) \sum_{y' \in \mathbf{Y}} H(x, y')\mathbb{P}_{\xi_0=\tilde{y}}\left\{\xi_1 = y'\right\} \\ &= \sum_{\tilde{y}' \in \mathbf{Y} \cup \{\wp\}} \tilde{H}(x, \tilde{y}')\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_1 = \tilde{y}'\right\} \\ &= \mathbb{E}_{\tilde{\xi}_0=\tilde{y}}\left[\tilde{H}(x, \tilde{\xi}_1)\right]. \end{aligned}$$

Therefore, η_t and $\tilde{\xi}_t$ are dual to one another with respect to $\tilde{H}$.

(b) Note first that by *i*, the set of linear combinations of $\tilde{H}(\cdot, \tilde{y})$, with $\tilde{y} \in \tilde{\mathbf{Y}}$, is dense on $C(\mathbf{X})$.

Denote by $\tilde{\Theta} := \{\theta \in \Theta : \ d(\theta) = 1\} \cup \{\wp\}$ the set of all absorbing states of $\tilde{\xi}_t$ and denote by τ the hitting time of $\tilde{\Theta}$ for $\tilde{\xi}_t$, that is,

$$\tau := \min\left\{s \geq 0 : \quad \tilde{\xi}_s \in \tilde{\Theta}\right\}.$$

From hypothesis *ii*, we obtain that there exists $a < 1$ such that for any non-absorbing state $\tilde{y}$ of $\tilde{\xi}_t$ we have that $d(\tilde{y}) \leq a$ and $\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_1 \notin \tilde{\Theta}\right\} \leq a$. It follows that for any $\tilde{y} \in \tilde{\mathbf{Y}} \setminus \tilde{\Theta}$, and any $s \geq 1$:

$$\begin{aligned}
\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_s \notin \tilde{\Theta}\right\} &= \sum_{\tilde{z}\in\tilde{\mathbf{Y}}\setminus\tilde{\Theta}} \mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_{s-1} = \tilde{z}\right\} \mathbb{P}_{\tilde{\xi}_{s-1}=\tilde{z}}\left\{\tilde{\xi}_s \notin \tilde{\Theta}\right\} \\
&\leq a \sum_{\tilde{z}\in\tilde{\mathbf{Y}}\setminus\tilde{\Theta}} \mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_{s-1} = \tilde{z}\right\} \\
&= a\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_{s-1} \notin \tilde{\Theta}\right\}.
\end{aligned}$$

Therefore, by induction and noting that $\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\{\tau = \infty\} \leq \mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_s \notin \tilde{\Theta}\right\}$ for any $s \geq 1$, we get $\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\{\tau = \infty\} \leq a^s$ and, since it holds for all $s \geq 1$, we deduce $\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\{\tau < \infty\} = 1$.

Defining, for any probability measure $\mu \in \mathcal{P}(\mathbf{X})$ and any $\tilde{y} \in \tilde{\mathbf{Y}}$, $\hat{\mu}(\tilde{y}) := \int_{\mathbf{X}} \tilde{H}(x,\tilde{y})d\mu(x)$, it follows that for any $\tilde{y} \in \tilde{\mathbf{Y}}$

$$\begin{aligned}
\lim_{s\to\infty} \hat{\mu}_s(\tilde{y}) &= \lim_{s\to\infty} \mathbb{E}_{\tilde{\xi}_0=\tilde{y}}\left[\hat{\mu}(\tilde{\xi}_s)\right] \\
&= \lim_{s\to\infty} \sum_{\tilde{\theta}\in\tilde{\Theta}} \mathbb{E}_{\tilde{\xi}_0=\tilde{y}}\left[\hat{\mu}(\tilde{\xi}_s)|\ \tilde{\xi}_\tau = \tilde{\theta},\ \tau \leq s\right] \mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_\tau = \tilde{\theta},\ \tau \leq s\right\} \\
&\quad + \lim_{s\to\infty} \mathbb{E}_{\tilde{\xi}_0=\tilde{y}}\left[\hat{\mu}(\tilde{\xi}_s)|\ \tau > s\right] \mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\{\tau > s\} \\
&= \sum_{\theta\in\Theta,\ d(\theta)=1} c(\theta)\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\left\{\tilde{\xi}_\tau = \theta\right\},
\end{aligned}$$

where the first equality follows from the fact that η_t and $\tilde{\xi}_t$ are dual to one another with respect to $\tilde{H}$ and equation (10), and the last one is due to *iii*, $\tilde{H}(\cdot, \wp) \equiv 0$ and $\tilde{\Theta} = \{\theta \in \Theta :\ d(\theta) = 1\} \cup \{\wp\}$.

□

Remark 2.2. *From the proof of Theorem 2.1, it is straightforward to check that condition* ii *can be replaced by* $\mathbb{P}_{\xi_0=y}\{\tau' < \infty\} = 1$ *for all* $y \in \mathbf{Y}$*, where* τ' *is the hitting time of* Θ *for* ξ_t*, since this implies* $\mathbb{P}_{\tilde{\xi}_0=\tilde{y}}\{\tau < \infty\} = 1$ *for all* $\tilde{y} \in \tilde{Y}$*.*

Remark 2.3. *As equation (7) can be seen as a discrete-time counterpart of (4), it is interesting to compare the form of the limit measure (11) with the expression given in (2.4) of [7] for continuous time:*

$$\hat{\nu}(y) = \mathbb{E}_{\xi_0=y}\left[e^{-\int_0^\tau V(\xi_s)ds}\mathbf{1}_{\{\tau<\infty\}}\right].$$

Note that, in [7], the expression is given through the dual process ξ_t, studying its complete evolution from zero time to the time of absorption, while here it is given only through the absorbing point of the modified process $\tilde{\xi}_t$ instead of the original dual process ξ_t.

Remark 2.4. *Theorem 2.1 gives conditions for the ergodicity of the process and, in that case, the form of the limit measure. Note that, even when the process is not ergodic, equation (7) and the computations made in the proof of the theorem can be used to study the behaviour of η_t as a function of the evolution of $\tilde{\xi}_t$.*

3. Nearest-neighbour probabilistic cellular automata

In this section we will use the duality equation (7) to study conditions for the ergodicity and the limit behaviour of some PCA models evolving on $\mathbb{Z}$.

First, let us introduce a definition:

Definition 3.1. *We say that a PCA η_t with state space $\mathbf{X} = W^{\mathbb{Z}} = \{1, \ldots, M\}^{\mathbb{Z}}$ is a nearest-neighbour PCA if $N = \{-1, 0, 1\}$.*

Thus, the evolution of such a process is given for any $i, j, k, m \in W$, $z \in \mathbb{Z}$ and $s \in \mathbb{N}$ by

$$\mathbb{P}\left\{\eta_{s+1}(z) = m \middle| \, \eta_s(z-1) = i, \eta_s(z) = j, \eta_s(z+1) = k\right\} = p_{ijk,m}.$$

We will treat two wide classes of nearest-neighbour PCA, including classical voter models, linear and non-linear voter models, noisy voter models, biased voter models and some competition models.

3.1. Multi-opinion voter models

Throughout this subsection, η_t will be a nearest-neighbour PCA with M states, and the state space of the dual processes will be $\mathbf{Y} = Y^{M-1}$ (where Y is the collection of finite subsets of $\mathbb{Z}$).

Let $H : \mathbf{X} \times \mathbf{Y} \to \mathbb{R}$ be the function defined for all $x = \{x(z) \in W\}_{z \in \mathbb{Z}} \in \mathbf{X}$ and $\mathbf{A} = (A_i)_{1 \le i \le M-1} \in \mathbf{Y}$ by

$$H(x, \mathbf{A}) = \begin{cases} 1 & , \quad \textit{if } x(z) = i, \ \forall z \in A_i, \ \forall i = 1, \ldots, M-1 \\ 0 & , \quad \textit{otherwise} \end{cases} . \tag{13}$$

Our aim is to construct a Markov chain $\mathbf{A}_t$ with state space $\mathbf{Y}$ and a function $d : \mathbf{Y} \to [0, \infty)$ that verify equation (7) with H defined in (13).

Suppose η_t and $\mathbf{A}_t = (A_{1,t}, \ldots, A_{M-1,t})$ are dual to one another with respect to (H, d), for some function d, and let $x \in \mathbf{X}$ and $\mathbf{A} = (A_i)_{1 \le i \le M-1} \in \mathbf{Y}$, where

$\{A_i\}_{1\le i\le M-1}$ is a pairwise disjoint family of subsets of $\mathbb{Z}$. Then, equation (7) can be written as:

$$\begin{aligned}\mathbb{P}_{\eta_0=x}\{\eta_1(z)=i,\forall z\in A_i,1\le i\le M-1\} &= d(\mathbf{A})\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\{x(z)=i,\forall z\in A_{i,1},1\le i\le M-1\}\\ &= d(\mathbf{A})\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{A_{i,1}\subseteq R_x^i,\ 1\le i\le M-1\right\},\end{aligned} \tag{14}$$

where $R_x^i:=\{z\in\mathbb{Z}:\ x(z)=i\}$. Note that given x and $\mathbf{A}$, then the left hand side can be written as the product

$$\begin{aligned}\mathbb{P}_{\eta_0=x}\{\eta_1(z)=i,\forall z\in A_i,1\le i\le M-1\} &= \prod_{i=1}^{M-1}\mathbb{P}_{\eta_0=x}\{\eta_1(z)=i,\forall z\in A_i\}\\ &= \prod_{i=1}^{M-1}\prod_{z\in A_i}\mathbb{P}_{\eta_0=x}\{\eta_1(z)=i\},\end{aligned}$$

as long as $A_1,\dots,A_{M-1}$ are pairwise disjoint. This factorization, due to the conditional independence of different particles, leads us to consider dual chains where the sets $A_{1,s},\dots,A_{M-1,s}$ evolve independently as long as they are disjoint; that is, given pairwise disjoint $A_1,\dots,A_{M-1}$, we require

$$\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\{A_{i,1}=B_i:\ 1\le i\le M-1\}=\prod_{i=1}^{M-1}\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\{A_{i,1}=B_i\}$$

for all pairwise disjoint $B_1,\dots,B_{M-1}\in\mathbf{Y}$. Indeed, we will consider processes in which the evolution of points belonging only to one component is independent of the rest. More precisely, for pairwise disjoint $A_{1,s},\dots,A_{M-1,s}$, a singleton $\{z\}$ contained in $A_{m,s}$ is substituted by $B_m(z)$, a subset of $\{z-1,z,z+1\}$, independently of the rest of singletons in $A_{m,s}$, with probabilities $\pi_m^{\cdot}$ given by



(15)

The set $A_{m,s+1}$ is the union of the resulting sets from the updating of all $\{z\}\subseteq A_{m,s}$, that is $A_{m,s+1}:=\bigcup_{z\in A_{m,s}}B_m(z)$. If the sets A_i are not pairwise disjoint, the point $\mathbf{A}=(A_1,\dots,A_{M-1})$ is absorbing for the dual process. Note that the evolution of this process can be seen as a multi-type coalescing branching process. Moreover, we will take the function d with the form

$$d(\mathbf{A}) = \prod_{i=1}^{M-1} d_i^{|A_i|}, \tag{16}$$

with $d_i \in [0, +\infty)$.

Now, let $x \in \mathbf{X}$ and $\mathbf{A} \in \mathbf{Y}$. If $A_1, \ldots, A_{M-1}$ are not pairwise disjoint, then both sides of (14) are zero. If $A_1, \ldots, A_{M-1}$ are pairwise disjoint, we have, due to the evolution of the dual, and noting that $R_x^1, \ldots, R_x^{M-1}$ are pairwise disjoint by definition,

$$\begin{aligned} d(\mathbf{A})\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{A_{i,1} \subseteq R_x^i :\ 1 \leq i \leq M-1\right\} = & \prod_{i=1}^{M-1} d_i^{|A_i|}\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{A_{i,1} \subseteq R_x^i\right\} \\ = & \prod_{i=1}^{M-1} d_i^{|A_i|} \prod_{\{z\}\subseteq A_i} \mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{B_i(z) \subseteq R_x^i\right\}, \end{aligned}$$

where $B_i(z)$ is the one-step evolution of $\{z\} \subseteq A_i$ in the dual process.

Therefore, equation (14) takes the form

$$\prod_{i=1}^{M-1} \prod_{z\in A_i} \mathbb{P}_{\eta_0=x}\left\{\eta_1(z) = i\right\} = \prod_{i=1}^{M-1} \prod_{\{z\}\subseteq A_i} d_i\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{B_i(z) \subseteq R_x^i\right\}.$$

Now, we look for conditions on the probabilities $\pi_m^{\cdot}$ such that η_t and $\mathbf{A}_t$ are dual with respect to (H, d). We must have $\mathbb{P}_{\eta_0=x}\left\{\eta_1(z) = m\right\} = d_m\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{B_m(z) \subseteq R_x^m\right\}$ for all $x \in \mathbf{X}$, $\mathbf{A} = (A_1, \ldots, A_{M-1})$ pairwise disjoint, $z \in \mathbb{Z}$ and $m = 1, \ldots, M-1$. These equations take the following form

$$\begin{array}{lcl} p_{ijk,m} & = & d_m\pi_m^{\emptyset} \\ p_{mjk,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{\ell}\right] \\ p_{imk,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{c}\right] \\ p_{ijm,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{r}\right] \\ p_{mmk,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{\ell} + \pi_m^{c} + \pi_m^{\ell c}\right] \\ p_{mjm,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{\ell} + \pi_m^{r} + \pi_m^{\ell r}\right] \\ p_{imm,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{c} + \pi_m^{r} + \pi_m^{cr}\right] \\ p_{mmm,m} & = & d_m\left[\pi_m^{\emptyset} + \pi_m^{\ell} + \pi_m^{c} + \pi_m^{r} + \pi_m^{\ell c} + \pi_m^{\ell r} + \pi_m^{cr} + \pi_m^{\ell cr}\right] \end{array}, \tag{17}$$

with $i, j, k \in W \setminus \{m\}$, together with

$$\pi_m^{\cdot} \geq 0, \qquad \pi_m^{\emptyset} + \pi_m^{\ell} + \pi_m^{c} + \pi_m^{r} + \pi_m^{\ell c} + \pi_m^{\ell r} + \pi_m^{cr} + \pi_m^{\ell cr} = 1. \tag{18}$$

Note that the last line of (17) is actually

$$p_{mmm,m} = d_m.$$

The existence of solutions for equations (17) and (18) requires extra assumptions on the process η_t. In fact, such equations may only have solutions if for any $m \in W \setminus \{M\}$ each of the parameters $p_{ijk,m}, p_{mjk,m}, p_{imk,m}, \ldots, p_{imm,m}$ does not depend on $i, j, k \neq m$.

That is to say that the probability of a site assuming a state m at time $s+1$ is a function of the positions of the state m in its neighbourhood at time s. Moreover, when (17) has a solution, we can note that $p_{mmm,m}=0$ implies $p_{ijk,m}=0$ for all $i,j,k\in W$, so the state m could not be taken by η_t. Hence, it is reasonable to assume $p_{mmm,m}>0$ for all $m\in W\setminus\{M\}$. Thus, $0<d_m\leq 1$ for all $m=1,\dots,M-1$.

Therefore, for any $m\in W\setminus\{M\}$ and $i,j,k\neq m$ we can denote $p_m:=p_{ijk,m}$, which can be interpreted as the probability that $\eta_1(z)$ spontaneously assumes the state m (i.e. when m is not present in the neighbourhood of the point). Thus, $p_{mmm,m}$ is equal to the probability that $\eta_1(z)$ does not spontaneously assume any state $k\neq m$, and for $m=1,\dots,M-1$, we have

$$d_m = 1-p_{mmm,M}-p+p_m$$

where $p:=\sum_{n=1}^{M-1}p_n$. Hence, (16) becomes

$$d(\mathbf{A})=\prod_{i=1}^{M-1}(1-p_{iii,M}-p+p_i)^{|A_i|}. \tag{19}$$

The equations (17) and (18) have solutions given by:

$$\begin{aligned}
\pi_m^{\emptyset} &= \frac{1}{d_m}p_m\\
\pi_m^{\ell} &= \frac{1}{d_m}(p_{mkk,m}-p_m)\\
\pi_m^{c} &= \frac{1}{d_m}(p_{kmk,m}-p_m)\\
\pi_m^{r} &= \frac{1}{d_m}(p_{kkm,m}-p_m)\\
\pi_m^{\ell c} &= \frac{1}{d_m}(p_{mmk,m}+p_m-p_{mkk,m}-p_{kmk,m})\\
\pi_m^{\ell r} &= \frac{1}{d_m}(p_{mkm,m}+p_m-p_{mkk,m}-p_{kkm,m})\\
\pi_m^{cr} &= \frac{1}{d_m}(p_{kmm,m}+p_m-p_{kmk,m}-p_{kkm,m})\\
\pi_m^{\ell cr} &= \frac{1}{d_m}(p_{mmm,m}+p_{mkk,m}+p_{kmk,m}+p_{kkm,m}-p_{kmm,m}-p_{mkm,m}-p_{mmk,m}-p_m)
\end{aligned} \tag{20}$$

with $k\neq m$, provided that

$$\begin{aligned}
&p_{mkk,m}\geq p_m\\
&p_{kmk,m}\geq p_m\\
&p_{kkm,m}\geq p_m\\
&p_{mmk,m}\geq p_{mkk,m}+p_{kmk,m}-p_m\\
&p_{mkm,m}\geq p_{mkk,m}+p_{kkm,m}-p_m\\
&p_{kmm,m}\geq p_{kmk,m}+p_{kkm,m}-p_m\\
&p_{mmm,m}+p_{mkk,m}+p_{kmk,m}+p_{kkm,m}\geq p_{kmm,m}+p_{mkm,m}+p_{mmk,m}+p_m
\end{aligned} \tag{21}$$

for all $m=1,\dots,M-1$, $k\in W\setminus\{m\}$. The inequalities (21) impose lower and upper bounds on the velocity for the growth of the probability that a site assumes the opinion m in the time $s+1$ because of the occurrence of opinions m in its neighbourhood in time s.

Thus, we have proved:

Theorem 3.2. *Let η_t be a nearest-neighbour PCA with state space $\mathbf{X}$, whose transition probabilities are such that for any $m = 1, \ldots, M-1$ each of the parameters $p_{ijk,m}, p_{mjk,m}, p_{imk,m}, \ldots, p_{imm,m}$ does not depend on $i, j, k \neq m$ and satisfy inequalities (21). Then there exist a Markov chain $\mathbf{A}_t$ with state space $\mathbf{Y}$, $H : \mathbf{X} \times \mathbf{Y} \to \mathbb{R}$, and $d : \mathbf{Y} \to (0, 1]$, such that η_t and $\mathbf{A}_t$ are dual to one another with respect to (H, d). Furthermore, if $p_{mmm,m} = 1$, for all $m = 1, \ldots, M-1$, then η_t and $\mathbf{A}_t$ are dual to one another with respect to H.*

$\square$

The following result follows from the application of Theorem 2.1 and Theorem 3.2.

Theorem 3.3. *Let η_t be a nearest-neighbour PCA whose transition probabilities are such that for any $m = 1, \ldots, M-1$ each of the parameters $p_{ijk,m}, p_{mjk,m}, p_{imk,m}, \ldots, p_{imm,m}$ does not depend on $i, j, k \neq m$ and satisfy inequalities (21), and let $\mathbf{A}_t = (A_{i,t})_{1 \leq i \leq M-1}$ be the process on $\mathbf{Y}$ defined by the transition probabilities (15) and (20). Then, the process is ergodic if and only if any of the following conditions holds:*

(i) *$p_{iii,M} > 0$ for all $i \in W \setminus \{M\}$, that is η_t admits spontaneous changes from any pure states $i \neq M$ to the state M;*

(ii) *$\exists m, n \in W \setminus \{M\}$, $m \neq n$, with $p_m, p_n > 0$, that is η_t admits spontaneous changes to at least two distinct states;*

(iii) *$\exists m \in W \setminus \{M\}$, with $p_m > 0$ and $p_{mmm,M} > 0$;*

(iv) *$\exists m \in W \setminus \{M\}$, with $p_m > 0$, and such that the process $\mathbf{A}_t$ starting at $\mathbf{A} = (\emptyset, \ldots, A_m, \ldots, \emptyset)$ has probability 1 of extinction for all $A_m \in Y$.*

Furthermore, when any of the above conditions holds, the unique equilibrium measure for η_t is defined for any $\mathbf{A} = (A_1, \ldots, A_{M-1}) \in \mathbf{Y}$, by

$$\hat{\nu}(\mathbf{A}) = \mathbb{P}_{\tilde{\mathbf{A}}_0 = \mathbf{A}} \left\{ \tilde{\mathbf{A}}_\tau = (\emptyset, \ldots, \emptyset) \right\},$$

where $\tilde{\mathbf{A}}_t$ is the process defined on $\mathbf{Y} \cup \{\wp\}$ with transition probabilities given in (12), and τ is the hitting time of $\{(\emptyset, \ldots, \emptyset)\} \cup \{\wp\}$ for $\tilde{\mathbf{A}}_t$.

Proof. We first check the sufficiency of the conditions for ergodicity.

With the previous construction, η_t and $\mathbf{A}_t$ are dual to one another with respect to (H, d). Also, by Lemma 3.1 of [7], the set of linear combinations of functions $H(\cdot, \mathbf{A})$ is dense in $C(\mathbf{X})$.

The set of absorbing states of $\mathbf{A}_t$ contains $(\emptyset, \ldots, \emptyset)$ together with all $\mathbf{A} = (A_1, \ldots, A_{M-1})$ which are not pairwise disjoint. Moreover, let us denote $\mathcal{A}$ the set of values $m \in \{1, \ldots, M-1\}$, if any, such that

$$p_m = p_{mkk,m} = p_{kkm,m} = p_{mkm,m} = 0, \qquad p_{kmk,m} = p_{mmk,m} = p_{kmm,m} = p_{mmm,m}.$$

Then, if $m \in \mathcal{A}$, we have $\pi^c_m = 1$ so the points in A_m do not change in time. Thus, the set Θ of absorbing points of $\mathbf{A}_t$ also includes the points $\mathbf{A} = (A_1, \ldots, A_{M-1})$ such that $A_i = \emptyset$ for all $i \notin \mathcal{A}$.

Let us see now that under any of the conditions *i*, *ii*, *iii* or *iv*, it holds $\{\theta \in \Theta : d(\theta) = 1\} = \{(\emptyset, \ldots, \emptyset)\}$. Obviously, $d((\emptyset, \ldots, \emptyset)) = 1$. Now, for $(A_1, \ldots, A_{M-1})$ not pairwise disjoint with $A_i \cap A_j \neq \emptyset$, we have, under any of *i*, *ii*, *iii* or *iv*, that at least one of $d_i = 1 - p_{iii,M} - p + p_i$ or $d_j = 1 - p_{jjj,M} - p + p_j$ must be smaller than 1, so from (19) we get $d(\mathbf{A}) < 1$. Last, let $i \in \mathcal{A}$; since $p_i = 0$ we have that $d_i = 1 - p_{iii,M} - p$; note that under *i* we have $p_{iii,M} > 0$, and under any of *ii*, *iii* or *iv*, there must be some $m \in W \setminus (\{M\} \cup \mathcal{A})$ with $p_m > 0$; therefore under any of *i*, *ii*, *iii* or *iv*, we get $d_i = 1 - p_{iii,M} - p \leq 1 - p_{iii,M} - p_m < 1$ and $d(\mathbf{A}) < 1$. Note that in particular, $\{\theta \in \Theta : d(\theta) = 1\} = \{(\emptyset, \ldots, \emptyset)\}$ implies that the condition *iii* of Theorem 2.1 holds with $c(\theta) = 1$.

Suppose *i* holds, therefore it follows $d(\mathbf{A}) \leq \max\limits_{k \in W \setminus \{M\}} \{1 - p_{kkk,M}\} < 1$ for any non-empty set $\mathbf{A} \in \mathbf{Y}$. If *ii* holds, then $d(\mathbf{A}) \leq \max\limits_{k=m,n} \{1 - p_k\} < 1$ for any non-empty set $\mathbf{A} \in \mathbf{Y}$. If *iii* holds, then for any non-empty set $\mathbf{A} \in \mathbf{Y}$ we have $d(\mathbf{A}) \leq \max\{1 - p_m,\ 1 - p_{mmm,M}\} < 1$. Hence, in all the cases, we apply Theorem 2.1 to conclude.

Now, suppose *iv* holds. We can consider that for any $k \in W \setminus \{M\}$, $k \neq m$, we have $p_k = 0$, and also $p_{mmm,M} = 0$, since if the contrary holds, we could conclude the ergodicity by *ii* or *iii*. Hence, for any $k \neq m$, it follows that $\pi^{\emptyset}_k = 0$, while $\pi^{\emptyset}_m > 0$ (see (20)). Notice that $\mathbf{Y} = \mathbf{Y}_1 \cup \mathbf{Y}_2$, where $\mathbf{Y}_1 := \{\mathbf{B} \in \mathbf{Y} :\ \exists k \neq m, B_k \neq \emptyset\}$ and $\mathbf{Y}_2 := \{\mathbf{B} \in \mathbf{Y} :\ \forall k \neq m, B_k = \emptyset\}$. For all $\mathbf{A} \in \mathbf{Y}_1$ we have $d(\mathbf{A}) \leq 1 - p_m < 1$, and if $\mathbf{A}_0 = \mathbf{A}$, then $\mathbf{A}_s \in \mathbf{Y}_1$ for all $s < \tau$. Thus, the process $\tilde{\mathbf{A}}_t$ starting in $\mathbf{A} \in \mathbf{Y}_1$ has probability 1 of being absorbed in $\wp$. On the other hand, for all $\mathbf{A} \in \mathbf{Y}_2$ we have $d(\mathbf{A}) = 1$ and if $\tilde{\mathbf{A}}_t$ starts in $\mathbf{A} \in \mathbf{Y}_2$, then we have $\tilde{\mathbf{A}}_s \in \mathbf{Y}_2$ for all $s < \tau$, and it coincides with the process $\mathbf{A}_t$ starting in $\mathbf{A} \in \mathbf{Y}_2$ which has probability 1 of being absorbed in $(\emptyset, \ldots, \emptyset)$. Therefore, in both cases, $\mathbb{P}_{\tilde{\mathbf{A}}_0 = \mathbf{A}} \{\tau < \infty\} = 1$. Thus, from the proof of Theorem 2.1 we conclude the ergodicity of η_t (see Remark 2.2).

Furthermore, in all the cases we get the expression of the equilibrium measure for η_t from Theorem 2.1.

For necessity we only have to show that the process is not ergodic in the two following cases: (a) $p_i = 0$ for all $i = 1, \ldots, M-1$ and there exists $j \neq M$ such that $p_{jjj,M} = 0$; or (b) $\exists! m \in \{1, \ldots, M-1\}$ with $p_m > 0$, $p_{mmm,M} = 0$, and there is some $A_m \in Y$ such that the process $\mathbf{A}_t$ starting in $\mathbf{A}_0 = (\emptyset, \ldots, A_m, \ldots, \emptyset)$ has probability strictly less than 1 of extinction. Case (a) is immediate since the probability measures δ_M and δ_j, concentrated on pure state configurations $x' \in \mathbf{X}$ and $x'' \in \mathbf{X}$, $x'(z) = M$ and $x''(z) = j$ for all $z \in \mathbb{Z}$, respectively, are both equilibrium measures for the process.

For case (b), note first that the measure δ_m is an equilibrium measure for the process. Now let $x \in \mathbf{X}$ with $x(z) \neq m$ for all $z \in \mathbb{Z}$ and $\mathbf{A} = (\emptyset, \ldots, A_m, \ldots, \emptyset)$ such that $\mathbf{A}_t$ has a probability less than 1 of extinction. Note that the set $\mathbf{Y}(m) :=$

$\{\mathbf{B} \in \mathbf{Y} : B_i = \emptyset, \forall i \neq m\}$ is closed for the evolution of the dual process (that is, $\mathbf{A}_t \in \mathbf{Y}(m)$ for all $t \geq 0$), and $d(\mathbf{B}) = 1$ for all $\mathbf{B} \in \mathbf{Y}(m)$. Then, for the process $\mathbf{A}_t$ enclosed in $\mathbf{Y}(m)$, the duality relation (7) takes the form of (2), and for the particular value of x and $\mathbf{A}$ we have

$$\begin{aligned}
\mathbb{P}_{\eta_0=x}\{\eta_s(z)=m,\ \forall z\in A_m\} &= \mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\{A_{m,s}\subseteq R_x^m\}\\
&= \mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\{A_{m,s}=\emptyset\}\\
&\leq \mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\{\tau<\infty\}<1.
\end{aligned}$$

Then, $\limsup_{s\to\infty}\mathbb{P}_{\eta_0=x}\{\eta_s(z)=m,\ \forall z\in A_m\}<1$ so the process starting with a measure with support in $\{x\in\mathbf{X}:\ x(z)\neq m, \forall z\in\mathbb{Z}\}$ is not converging to δ_m and the process is not ergodic.

□

Remark 3.4. *Theorems 3.2 and 3.3 can be extended to a wide class of PCA with finite range interaction on the integer lattice* $\mathbb{Z}^d$ *(1), which are such that* $\mathbb{P}\{\eta_{s+1}(z)=w|\ \eta_s(z+n)=w_n,\ \forall n\in N\}$ *does not depend on* $w_n\neq w$. *In fact, for any* d *and* $N\subset\mathbb{Z}^d$ *we can define a dual process* ξ_t *on* $\mathbf{Y}=Y^{M-1}$, *where* Y *is the collection of finite subsets of* $\mathbb{Z}^d$, *and a point* z *belonging to* $A_{m,s}$ *is substituted by a subset of* $\{z+n:\ n\in N\}$. *Thus, in an analogous way to that in (21) it is possible to find inequalities that delimit the class of PCA for which we can extend our results.*

The PCA model presented in this subsection recover many one-dimensional models that have been referred in the literature. In fact, our model recovers M-state linear and non-linear PCA (which were studied in the 2-state version by [13] and [14]), presenting new results for them.

Example 3.5. (Multi-opinion noisy voter model) *Suppose* η_t *is defined as follows: For each* $m\in W=\{1,\ldots,M\}$, *take* $q_m\in[0,1]$, *and define* $q=\sum_{m\in W}q_m$; *set* $\alpha,\beta,\gamma\geq 0$, *such that* $\alpha+\beta+\gamma=1-q$; *finally suppose* η_t *has transition probabilities given by*

$$p_{ijk,m} := \alpha\mathbf{1}_{\{i\}}(m)+\beta\mathbf{1}_{\{j\}}(m)+\gamma\mathbf{1}_{\{k\}}(m)+q_m.$$

Note that for any $m\in W$ *and* $i,j,k\in W\setminus\{m\}$ *we have* $q_m=p_{ijk,m}=:p_m$ *(which here is also defined for* $m=M$*). Using the language of voter models,* q_m *is the probability of a voter assuming the opinion m, independently of her own opinion, as well as the opinions of her neighbours in the previous step, while* α, β *and* γ *are the weights given by each voter to her own opinion and her neighbours' opinions.*

Corollary 3.6. *Let* η_t *be a multi-opinion noisy voter model. Then there exists a coalescing random walk* $\mathbf{A}_t$ *on the family of all finite subsets of* $\mathbb{Z}$ *such that* η_t *and* $\mathbf{A}_t$ *are dual to one another with respect to* (H,d) *defined in (13) and (19). Moreover,* η_t *is ergodic if and only if there exists* $m\in\{1,\ldots,M\}$ *such that* $p_m>0$.

Proof. It is easy to check that for any $m = 1, \ldots, M-1$ the parameters $p_{ijk,m}, p_{mjk,m}$, $p_{imk,m}, \ldots, p_{imm,m}$ do not depend on $i, j, k \neq m$ and satisfy inequalities (21). Hence, we can apply Theorem 3.2 to deduce the existence of a dual with respect to (H, d), where H is given by (13) and d is given by (19).

For the second part of the result, if $p_m = 0$ for all m, then any probability measure concentrated on a pure state configuration $x \in \mathbf{X}$, $x(z) = m$ for all $m \in \mathbb{Z}$ and any $m \in W$, is an equilibrium measure. For sufficiency, note that (21) holds regardless of the order of the opinions, and thus we can set M to have $p_M > 0$ and ergodicity follows from Theorem 3.3 *i*.

□

Example 3.7. (Domany-Kinzel model) *If we consider the case* $W = \{0, 1\}$ *and such that the probability of a site* z *assuming* 1 *at time* $s+1$ *is a function of the number of* 1*s in the neighbourhood* $\{z-1, z+1\}$ *at time* s*, then we have the Domany-Kinzel model (6) with parameters*

$$\begin{aligned} a_0 &:= p_{000,1} = p_{010,1}; \\ a_1 &:= p_{100,1} = p_{110,1} = p_{011,1} = p_{001,1}; \\ a_2 &:= p_{101,1} = p_{111,1}. \end{aligned}$$

We recall that the Domany-Kinzel model was introduced in [15] and has been extensively studied (see for instance [12], [16], [17] and [18]) due to its useful applicability in percolation theory and phase-transition theory among others.

Corollary 3.8. *Let* η_t *be the Domany-Kinzel model (6) with parameters* $0 \leq a_0 \leq a_1 \leq a_2 \leq 1$*. Then there exists a coalescing random walk* $\mathbf{A}_t$ *on the family of all finite subsets of* $\mathbb{Z}$*, such that* η_t *and* $\mathbf{A}_t$ *are dual to one another with respect to the pair* (H, d) *defined by (13) and (19). Furthermore, if one of the following conditions holds*

(i) $a_0 + a_2 \geq 2a_1$ *and* $a_2 < 1$*;*

(ii) $a_0 + a_2 < 2a_1$ *and* $a_0 > 0$*;*

(iii) $a_0 > 0$ *and* $a_2 < 1$*;*

(iv) $a_0 = 0$, $a_1 < 1/2$ *and* $a_2 < 1$*;*

(v) $a_0 > 0$, $a_1 > 1/2$ *and* $a_2 = 1$*;*

then η_t *is ergodic.*

Proof. If $a_1 \leq \frac{1}{2}(a_0 + a_2)$, then η_t satisfies all hypotheses of Theorem 3.2 (identifying the state 0 here with the the state M in Theorem 3.2). Hence, we have that $\mathbf{A}_t$ is defined by the transition probabilities $\pi^{\emptyset} := a_0/a_2$, $\pi^{\ell} = \pi^{r} := (a_1 - a_0)/a_2$, and $\pi^{\ell r} := (a_2 - 2a_1 + a_0)/a_2$.

On the other hand, if $a_1 > \frac{1}{2}(a_0 + a_2)$, then we can consider the description of the process by the probabilities of the occurrence of 0, that is the Domany-Kinzel model with parameters

$$a_i^* := 1 - a_{2-i}, \qquad i = 0, 1, 2.$$

It is easy to see that $0 \leq a_0^* \leq a_1^* \leq a_2^* \leq 1$, and $a_1^* < \frac{1}{2}(a_0^* + a_2^*)$. Therefore, we can again apply Theorem 3.2 (now the state 1 plays the role of the state M in Theorem 3.2) and find $\mathbf{A}_t^*$, which is defined by the transition probabilities $\pi_*^{\emptyset} := a_0^*/a_2^*$, $\pi_*^{\ell} = \pi_*^{r} := (a_1^* - a_0^*)/a_2^*$, and $\pi_*^{\ell r} := (a_2^* - 2a_1^* + a_0^*)/a_2^*$.

Furthermore, we can obtain sufficient conditions for ergodicity:

(i) Identifying the state M with 0 in Theorem 3.2 and $p_{111,0} = 1 - a_2 > 0$, the result follows from Theorem 3.3 *i*.

(ii) We now identify M with 1 so $p_{000,1} = 1 - a_2^* = a_0 > 0$ and the conclusion follows from Theorem 3.3 *i*.

(iii) We identify M with 0 or 1 depending on $(a_2 + a_0)/2$ being greater or smaller than a_1. In both cases, for $i \neq M$ we have $p_{iii,M} > 0$ and the result follows from Theorem 3.3 *i*.

(iv) Suppose $a_0 = 0$, $a_1 < 1/2$, and $a_2 < 1$. We have two possible cases:

   (a) If $a_1 \leq a_2/2$, then we can use the dual process $\mathbf{A}_t$ and, since $p_{111,0} = 1 - a_2 > 0$ and the state 0 plays the role of the state M in the Theorem 3.2, we can deduce the ergodicity of η_t from Theorem 3.3.*i*.

   (b) If $a_1 > a_2/2$, then consider the dual process $\mathbf{A}_t^*$ (that is, with $M = 1$, $p_{000,1} = 0$, $p_0 = 1 - a_2 > 0$), starting from any $\mathbf{A}_0 \in \mathbf{Y}$. Let us show that $\mathbf{A}_t^*$ has probability 1 of extinction.
   Consider the branching process χ_t on $\mathbb{Z}$ defined as follows: Let χ_s be the number of individuals in the sth generation of some population; assume each individual independently will give rise to 0 offsprings with probability $\pi_*^{\emptyset}$, 1 offspring with probability $(\pi_*^{\ell} + \pi_*^{r})$, or 2 offsprings with probability $\pi_*^{\ell r}$; then χ_{s+1} is the number of individuals in the next generation, that is the total number of offsprings generated by the individuals in the generation s. Since $a_1 < 1/2$, it follows that the expected number of offsprings of each individual is $(\pi_*^{l} + \pi_*^{r}) + 2\pi_*^{lr} = 2a_1 < 1$, which implies the branching process χ_t vanishes with probability 1 (see [19]). Identifying χ_t starting with $|\mathbf{A}_0|$ individuals as an upper bound on the number of individuals of $\mathbf{A}_t^*$, we have that the dual process $\mathbf{A}_t^*$ has probability 1 of being absorbed in $\emptyset$ and we can use Theorem 3.3.*iv* to deduce the ergodicity of η_t.

(v) If $a_0 > 0$, $a_1 > 1/2$ and $a_2 = 1$, then $a_0^* = 0$, $a_1^* < 1/2$ and $a_2^* < 1$, and therefore we get the result reasoning as in *iv*.

□

Remark 3.9. *As we said in the Introduction, the inclusion of the function d in our definition of duality (7) allows, in some cases, to analyze processes which do not have a dual with respect to a given function H in terms of the classical duality equation (2). In fact, from the construction of Theorem 3.2, we can see that if we consider the classical duality (2) with respect to the function H defined in (13) we would obtain a system of equations similar to (17), but without the factor d_m multiplying the right side of the equations. Therefore, for the existence of a solution of the system, we would need the extra hypothesis $p_{mmm,m} = 1$ for all $m = 1, \dots, M-1$. In particular, $M \geq 3$ would imply that every state $m = 1, \dots, M-1$ should have $p_m = 0$ and $p_{mmm,M} = 0$; while if $M = 2$, we could have $p_1 > 0$ but not $p_{111,2} > 0$. Note that the additional hypothesis of $p_{mmm,m} = 1$ for all $m \neq M$ would imply that if $M \geq 3$, then η_t is not ergodic; and if $M = 2$, then η_t would be ergodic if and only if p_1 was large enough in order for condition of Theorem 3.3* iii *to hold.*

In Example 3.5, if we had used the classical duality equation (2), we would be restricted to consider models with every $q_m = 0$ (if $M \geq 3$) or with only one $q_m \neq 0$ (if $M = 2$). On the other hand, in Example 3.7, and due to symmetry of states '0' and '1' in the Domany-Kinzel model, we could have either $a_0 > 0$ or $a_2 > 0$, but not both a_0 and a_2 positive.

3.2. Multi-state monotone models

In the previous subsection we showed that a necessary condition for a process to have a dual with respect to (H, d) defined in (13) and (19) is that for any $m = 1, \dots, M-1$ each of the parameters $p_{\dots,m}$ depends only on the presence (and position) of the value m in the neighbourhood of the point. However, there are models (for instance, when the set W has an order) where not only the presence (and position) of the value m is important for the probability of changing to m but also the presence (and position) of values bigger or smaller than m is. In order to find a dual for these processes, we will use a duality function H different to (13) and allow more transitions for the dual than we did in Subsection 3.1; as we will see, in this case, the evolution of any two subsets A_i and A_j has a greater interdependence than in the non-biased case.

For simplicity, we consider only the case of η_t being a nearest-neighbour PCA for which the probability of a fixed site assuming some state at time $s+1$ does not depend on its own state at time s. That is, we assume that

$$p_{ik,m} := p_{ijk,m}, \qquad \forall i, j, k, m \in W. \tag{22}$$

Throughout this subsection, the state space of the dual process will be $\mathbf{Y} = \{\mathbf{A} \in Y^{M-1} : A_1 \subseteq \cdots \subseteq A_{M-1}\}$ and the function $H : \mathbf{X} \times \mathbf{Y} \to \mathbb{R}$, defined by

$$H(x, \mathbf{A}) = \begin{cases} 1 & , \quad if\ x(z) \leq k,\ \forall z \in A_k,\ \forall k = 1, \dots, M-1 \\ 0 & , \quad otherwise \end{cases}. \tag{23}$$

The class of dual processes we consider is composed of processes $\mathbf{A}_t (= \xi_t)$ starting at $\mathbf{A}_0 \in \mathbf{Y}$ whose transition probabilities from $\mathbf{A}_s = (A_{i,s})_{1 \leq i \leq M-1}$ to $\mathbf{A}_{s+1} =$

$(A_{i,s+1})_{1\le i\le M-1}$ are defined as follows. Let $(\emptyset,\dots,\emptyset)$ be an absorbing state for $\mathbf{A}_t$ and, under the convention that $A_{0,s}=\emptyset$, assume that for any $k,m,n\in\{1,\dots,M-1\}$ each singleton $\{z\}\in A_{k,s}\backslash A_{k-1,s}$ will generate an element independently of other $z'\in A_{M-1,s}$, $\mathbf{B}(z)=(B_i(z))_{1\le i\le M-1}\in\mathbf{Y}$ as follows:

- with probability $\pi_k^{\emptyset}$
$$B_i(z)=\emptyset, \qquad 1\le i\le M-1;$$
- with probability $\pi_{k,m}^{\ell}$
$$B_i(z)=\begin{cases}\emptyset & , \; if \quad 1\le i<m\\ \{z-1\} & , \; if \quad m\le i\le M-1;\end{cases}$$
- with probability $\pi_{k,m}^{r}$
$$B_i(z)=\begin{cases}\emptyset & , \; if \quad 1\le i<m\\ \{z+1\} & , \; if \quad m\le i\le M-1;\end{cases}$$
- with probability $\pi_{k,mn}$
$$B_i(z)=\begin{cases}\emptyset & , \; if \quad 1\le i<\min\{m,n\}\\ \{z-1\} & , \; if \quad m\le i<n\\ \{z+1\} & , \; if \quad n\le i<m\\ \{z-1,z+1\} & , \; if \quad \max\{m,n\}\le i\le M-1;\end{cases}$$

with
$$\pi_k^{\emptyset}+\sum_{m=1}^{M-1}\pi_{k,m}^{\ell}+\sum_{n=1}^{M-1}\pi_{k,n}^{r}+\sum_{m=1}^{M-1}\sum_{n=1}^{M-1}\pi_{k,mn}=1.$$

We define $\mathbf{A}_{s+1}=(A_{i,s+1})_{1\le i\le M-1}$, where
$$A_{i,s+1}=\bigcup_{z\in A_{M-1}}B_i(z).$$

Note that $\mathbf{A}_{s+1}$ also belongs to $\mathbf{Y}$.

Equation (7) is, in this case, for each $x\in\mathbf{X}$, $\mathbf{A}\in\mathbf{Y}$,

$$\mathbb{P}_{\eta_0=x}\left\{\eta_1(z)\le k:\forall z\in A_k, 1\le k\le M-1\right\}=d(\mathbf{A})\mathbb{P}_{\mathbf{A}_0=\mathbf{A}}\left\{A_{k,1}\subseteq\mathcal{R}_x^k:1\le k\le M-1\right\}, \tag{24}$$

where $\mathcal{R}_x^k:=\{z\in\mathbb{Z}:\ x(z)\le k\}$.

The left hand side of (24) takes the form:

$$\prod_{k=1}^{M-1}\prod_{z\in A_k\backslash A_{k-1}}\mathbb{P}_{\eta_0=x}\left\{\eta_1(z)\le k\right\}$$

and, assuming that $d : \mathbf{Y} \to \mathbb{R}$ can be written as

$$d(\mathbf{A}) = \prod_{k=1}^{M-1} d_k^{|A_k \setminus A_{k-1}|}$$

for some $d_1, \ldots, d_{M-1} \in [0,1]$ (the product over the empty set is taken as 1), the right hand side of (24) is

$$\prod_{k=1}^{M-1} \prod_{\{z\} \subseteq A_k \setminus A_{k-1}} d_k \mathbb{P}_{\mathbf{A}_0 = \mathbf{A}} \left\{ B_m(z) \subseteq \mathcal{R}_x^m : 1 \le m \le M-1 \right\}.$$

Then, we must find solutions for the set of equations

$$\mathbb{P}_{\eta_0 = x} \left\{ \eta_1(z) \le k \right\} = d_k \mathbb{P}_{\mathbf{A}_0 = \mathbf{A}} \left\{ B_m(z) \subseteq \mathcal{R}_x^m : 1 \le m \le M-1 \right\}, \tag{25}$$

for all $x \in \mathbf{X}$, $\mathbf{A} \in \mathbf{Y}$, $k = 1, \ldots, M-1$ and $z \in A_k \setminus A_{k-1}$. Writing $S_{i,j}^k = \sum_{m=1}^k p_{ij,m}$ and noting that, for $1 \le i,j \le M$, $x(z-1) = i$, $x(z+1) = j$, we have

$$\mathcal{R}_x^m \supseteq \begin{cases} \{z-1\} & , \quad if\ i \le m < j \\ \{z+1\} & , \quad if\ j \le m < i \\ \{z-1, z+1\} & , \quad for\ m \ge i, j \end{cases}.$$

The equations (25) are

$$\begin{aligned}
S_{M,M}^k &= d_k \pi_k^{\emptyset} \\
S_{M-1,M}^k &= d_k \left[\pi_k^{\emptyset} + \pi_{k,M-1}^{\ell} \right] \\
S_{M,M-1}^k &= d_k \left[\pi_k^{\emptyset} + \pi_{k,M-1}^{r} \right] \\
S_{M-1,M-1}^k &= d_k \left[\pi_k^{\emptyset} + \pi_{k,M-1}^{\ell} + \pi_{k,M-1}^{r} + \pi_{k,M-1\ M-1} \right] \\
S_{M-2,M-1}^k &= d_k \left[\pi_k^{\emptyset} + \pi_{k,M-1}^{\ell} + \pi_{k,M-2}^{\ell} + \pi_{k,M-1}^{r} + \pi_{k,M-1\ M-1} + \pi_{k,M-2\ M-1} \right] \\
&\vdots \\
S_{i,j}^k &= d_k \left[\pi_k^{\emptyset} + \sum_{m=i}^{M-1} \pi_{k,m}^{\ell} + \sum_{n=j}^{M-1} \pi_{k,n}^{r} + \sum_{m=i}^{M-1} \sum_{n=j}^{M-1} \pi_{k,mn} \right] \\
&\vdots \\
S_{1,1}^k &= d_k \left[\pi_k^{\emptyset} + \sum_{m=1}^{M-1} \pi_{k,m}^{\ell} + \sum_{n=1}^{M-1} \pi_{k,n}^{r} + \sum_{m=1}^{M-1} \sum_{n=1}^{M-1} \pi_{k,mn} \right]
\end{aligned} \tag{26}$$

for $k = 1, \dots, M-1$. Due to the last line of the above equations, we deduce that $d_k = S^k_{1,1}$ and then

$$d(\mathbf{A}) = \prod_{k=1}^{M-1} \left(\sum_{m=1}^{k} p_{11,m} \right)^{|A_k \setminus A_{k-1}|} . \tag{27}$$

We assume $p_{11,1} = S^1_{1,1} > 0$, for otherwise, under (26), $S^1_{i,j} = 0$ for all $i, j \in \{1, \dots, M\}$ and opinion 1 cannot be taken by the process. Therefore, $d_k = S^k_{1,1} > 0$ for all $k \in \{1, \dots, M-1\}$ and the solution of (26) is given by:

$$\begin{aligned}
\pi^{\emptyset}_k &= \frac{S^k_{M,M}}{d_k} \\
\pi^{\ell}_{k,m} &= \frac{S^k_{m,M} - S^k_{m+1,M}}{d_k} \\
\pi^{r}_{k,m} &= \frac{S^k_{M,m} - S^k_{M,m+1}}{d_k} \\
\pi_{k,mn} &= \frac{S^k_{m,n} - S^k_{m+1,n} - S^k_{m,n+1} + S^k_{m+1,n+1}}{d_k}
\end{aligned} \tag{28}$$

since

$$\begin{aligned}
d_k \sum_{m=i}^{M-1} \pi^{\ell}_{k,m} &= \sum_{m=i}^{M-1} \left(S^k_{m,M} - S^k_{m+1,M} \right) = S^k_{i,M} - S^k_{M,M}, \\
d_k \sum_{n=j}^{M-1} \pi^{r}_{k,n} &= \sum_{n=j}^{M-1} \left(S^k_{M,n} - S^k_{M,n+1} \right) = S^k_{M,j} - S^k_{M,M}, \\
d_k \sum_{m=i}^{M-1} \sum_{n=j}^{M-1} \pi_{k,mn} &= \sum_{m=i}^{M-1} \sum_{n=j}^{M-1} \left(S^k_{m,n} - S^k_{m+1,n} - S^k_{m,n+1} + S^k_{m+1,n+1} \right) \\
&= \sum_{m=i}^{M-1} \left(S^k_{m,j} - S^k_{m,M} - S^k_{m+1,j} + S^k_{m+1,M} \right) \\
&= S^k_{i,j} - S^k_{M,j} - S^k_{i,M} + S^k_{M,M}.
\end{aligned} \tag{29}$$

In order to have the nonnegativity of (28), we need

$$S^k_{m,M} \geq S^k_{m+1,M}, \qquad S^k_{M,n} \geq S^k_{M,n+1}, \tag{30}$$

$$S^k_{m,n} - S^k_{m+1,n} - S^k_{m,n+1} + S^k_{m+1,n+1} \geq 0, \tag{31}$$

for all $1 \leq m, n \leq M-1$. Note that (30) and (31) together imply $S^k_{i,j} \geq S^k_{m,n}$ for all $1 \leq i \leq m \leq M$, $1 \leq j \leq n \leq M$ and $k = 1, \dots, M$, that is

$$\sum_{\ell=1}^{k} p_{ij,\ell} \geq \sum_{\ell=1}^{k} p_{mn,\ell}$$

which means that the process η_t is **monotone** with respect to the coordinatewise order in $\mathbf{X}$ induced by the total order $1 \leq 2 \leq \cdots \leq M$ in W (see [21]). Therefore, condition (30) can be substituted by the monotonicity of the process.

On the other hand, conditions (30) and (31) can be written equivalently as saying that $S^k_{i,j} = 1 - F^k(i,j)$ for each $k = 1, \dots M-1$, where F^k is a distribution function on $\mathbb{R}^2$.

We have proved the following result:

Theorem 3.10. *Let η_t be a nearest-neighbour PCA with state space $\mathbf{X}$ whose transition probabilities satisfy (22). Then, there exist a dual process $\mathbf{A}_t$ with state space $\mathbf{Y}$ such that η_t and $\mathbf{A}_t$ are dual to one another with respect to (H, d) defined in (23) and (27) if and only if η_t is monotone and its transition probabilities satisfy (31). Furthermore, if $p_{11,1} = 1$, then η_t and $\mathbf{A}_t$ are dual to one another with respect to H.*

□

Corollary 3.11. *A monotone nearest-neighbour PCA whose transition probabilities satisfy (31) is ergodic if $p_{11,M} > 0$. In such case, the unique equilibrium measure for η_t is defined for any $\mathbf{A} = (A_1, \dots, A_{M-1}) \in \mathbf{Y}$, by*

$$\hat{\nu}(\mathbf{A}) = \mathbb{P}_{\tilde{\mathbf{A}}_0 = \mathbf{A}} \left\{ \tilde{\mathbf{A}}_\tau = (\emptyset, \dots, \emptyset) \right\},$$

where $\tilde{\mathbf{A}}_t$ is the process defined on $\mathbf{Y} \cup \{\wp\}$ with transition probabilities given in (28), and τ is the hitting time of $\{(\emptyset, \dots, \emptyset)\} \cup \{\wp\}$ for $\tilde{\mathbf{A}}_t$.

Proof. By Theorem 3.10 there exist a Markov chain $\mathbf{A}_t$ and a pair of functions (H, d), such that η_t and $\mathbf{A}_t$ are dual to one another with respect to (H, d).

We now apply Theorem 2.1 to get the result. In fact, by Lemma 3.2 of [7] the set of linear combinations of the functions $H(\cdot, \mathbf{A})$ with $\mathbf{A} \in \mathbf{Y}$ is dense in $C(\mathbf{X})$. Also, $(\emptyset, \dots, \emptyset)$ is the unique absorbing state θ of $\mathbf{A}_t$ such that $d(\theta) = 1$, and $H(\cdot, (\emptyset, \dots, \emptyset)) \equiv 1$. Moreover, for any $\mathbf{A} \neq (\emptyset, \dots, \emptyset)$, $d(\mathbf{A}) = \prod_{k=1}^{M-1} (1 - \sum_{j=k+1}^{M} p_{11,j})^{|A_k \setminus A_{k-1}|} \leq 1 - p_{11,M}$. Thus, if $p_{11,m} > 0$, hypotheses *ii* and *iii* (with $c(\emptyset) = 1$) of Theorem 2.1 hold and the result is proved.

□

Remark 3.12. *Note that as in the case of the multi-opinion voter models, we can extend Theorem 3.10 and Corollary 3.11 to a wide class of multi-state monotone biased models with state space $W^{\mathbb{Z}^d}$ and finite range interaction.*

We notice that the processes η_t that verify the conditions of Theorem 3.10 include many PCA models, such as multi-opinion noisy biased voter models and multi-species competition models.

Example 3.13. (Multi-species competition model) *Suppose M distinct species live in the space $\mathbb{Z}$. Each site of the space contains only one individual. The spread mechanism for all species is the same: each individual produces one offspring (independently of other individuals) that will leave to colonize a neighbouring site, killing the old particle that occupied the site and competing against the offspring of others individuals. Moreover, we include the possibility of a spontaneous appearance of an individual of any type (for instance, coming from outside).*

We consider such a model described as follows: Suppose η_t is a nearest-neighbour process with M states such that the probability $p_{ij,m}$ that a site assumes the state m in the next step when its left neighbour is in the state i and its right neighbour is in the state j is given for any $j, k \neq m$ and $i < m$ by

$$p_{jk,m} = p_m$$

$$p_{im,m} = p_m + (1-p)\alpha_m$$

$$p_{im,i} = p_i + (1-p)(1-\alpha_m)$$

$$p_{mi,m} = p_m + (1-p)\beta_m$$

$$p_{mi,i} = p_i + (1-p)(1-\beta_m)$$

$$p_{mm,m} = 1 - p + p_m,$$

where $p = \sum_{m=1}^{M} p_m$, $\alpha_M \geq \alpha_{M-1} \ldots \geq \alpha_2 \geq 1/2$ and $\beta_M \geq \beta_{M-1} \ldots \geq \beta_2 \geq 1/2$.

The parameters α_m and β_m can be interpreted as the competitiveness of species m with relation to species $i < m$, and p_m corresponds to probabilities that spontaneous mutations to m occur. Note that the order in the αs and βs imply that species are ordered from weakest 1 to strongest M. Usually, we have $\alpha_m = \beta_m$ for all m, which means there is not a directional component in the competition.

Competition models belong to the class of biased voter models. The biased voter model was introduced in the 2-state version by Williams and Bjerknes [20] as a model of skin cancer, and has been applied to study several phenomena (for instance, see [22], [23], and [24]).

Since

$$S^k_{i,j} = \sum_{\ell=1}^{k} p_\ell + (1-p)\left((1-\alpha_j)\mathbf{1}_{\{i \leq k < j\}} + (1-\beta_i)\mathbf{1}_{\{j \leq k < i\}} + \mathbf{1}_{\{k \geq i,j\}}\right),$$

it is not hard to check that η_t is monotone and (31) holds for it. Therefore, by Corollary 3.11, if η_t admits a spontaneous change to the state M, then it is ergodic.

Example 3.14. *Consider the case of monotone processes for which* $S^k_{i,j} = g_k(i+j)$, *where* $g_k : \{2, \dots, 2M\} \to [0,1]$. *In such case, writing* $l := i+j$, *condition (31) takes the form*

$$\frac{1}{2}\left(g_k(l) + g_k(l+2)\right) \geq g_k(l+1),$$

which means that (31) is satisfied if and only if the functions g_k *are convex. In this case, we have that* η_t *is ergodic if* $g_{M-1}(2) < 1$. *In fact, if* $g_{M-1}(2) < 1$, *then* $p_{11,M} = 1 - S^{M-1}_{1,1} = 1 - g_{M-1}(2) > 0$ *and we can use Corollary 3.11 to deduce the ergodicity of the process.*

Remark 3.15. *As in Subsection 3.1, our definition (7) of duality with respect to* (H, d) *allows us to include processes that do not have a dual in terms of the classical definition (2). In fact, we could not obtain a version of Theorem 3.10 for classical duality (2) with respect to the function* H *defined in (23) unless we imposed another restrictive condition on* η_t *since we would obtain a system of equations similar to (26), but without the factor* d_k *multiplying the right side of each equation. Therefore, we should have* $S^k_{1,1} = 1$ *for any* $k = 1 \dots, M-1$, *which is equivalent to imposing the condition* $p_{11,1} = 1$. *In such case, we would have* $p_{11,M} = 0$ *and it would not be possible to obtain a sufficient condition for ergodicity in the form of Corollary 3.11.*

In particular, in Example 3.13, classical duality (2) with respect to H *would need* $p_m = 0$ *for any* $m \neq 1$ *(which implies* $p_M = 0$, *and therefore we could not deduce the ergodicity), while in Example 3.14 we should impose the condition* $g_1(2) = 1$ *(which implies* $g_{M-1}(2) = 1$*) and therefore we could not deduce the ergodicity either.*

4. Concluding remarks

In this work, we have presented a new duality equation (7) and shown its usefulness in the study of PCA. We believe that there is still room for future research on this topic. In particular we are considering the following issues:

(i) Studying in more detail the equilibrium measures (proportion of values, spatial correlations, etc.) when the processes are ergodic in Examples 3.5, 3.7 and 3.13.

(ii) In the case of processes having a dual with respect to (H, d) in Theorems 3.2 and 3.10 but not being ergodic, studying their asymptotic behaviour using their dual processes.

(iii) Studying other classes of functions H giving conditions on the transition probabilities for the existence of a dual with respect to (H, d).

Acknowledgments

We thank the referees for their careful reading and useful comments.

This work was supported by projects MTM2004-01175 and MTM2007-63769 of MEC, and research group “Modelos Estocásticos” (DGA). M. Sobottka was partially supported by Nucleus Millennium Information and Randomness ICM P04-069-F and project DIUC 207.013.030-1.0.

References

[1] Lebowitz J. L., Maes C. and Speer E. R., *Statistical mechanics of probabilistic cellular automata*, 1990 *J. of Statistical Physics* **59** 117-170.

[2] Toom A. L., Vasilyev N. B., Stavskaya O. N., Mityushin L. G., Kurdyumov G. L. and Pirogov S. A., *Discrete local Markov systems*, 1990 in:*R. L. Dobrushin, V. I. Kryukov, A. L. Toom, eds., Stochastic Cellular Systems: Ergodicity, Memory, Morphogenesis* 1-182.

[3] Liggett T. M., *Interacting particle systems*, 1985 Springer-Verlag, New York.

[4] Durrett R., *Lectures Notes on Particle Systems and Percolation*, 1988 Wadsworth, Belmont, CA.

[5] Gray L., *Duality for general attractive spin systems with applications in one dimension*, 1986 *Ann. Prob.* **14** 371-396.

[6] Katori M., *Reformulation of Gray's duality for attractive spin systems and its applications*, 1994 *J. Phys. A* **27** 3191-3211.

[7] López F. J. and Sanz G., *Duality for general interacting particle systems*, 2000 *Markov Processes and Relat. Fields* **6** 305-328.

[8] Katori M., Konno N., Sudbury A. and Tanemura H., *Dualities for the Domany-Kinzel model*, 2004 *J. of Theoretical Probability* **17** 131-144.

[9] Sudbury A. and Lloyd P., *Quantum operators in classical probability theory. II. The concept of duality in interacting particle systems*, 1995 *Ann. Prob.* **23** 1816-1830.

[10] Konno N., *Dualities for a class of finite range probabilistic cellular automata in one dimension*, 2002 *J. of Statistical Physics* **106** 915-922.

[11] Konno N. and Ma X., *Site-bond representation and self-duality for totalistic probabilistic cellular automata in one dimension*, 2005 *J. of Statistical Mechanics: Theory and Experiment* P03002.

[12] Konno N., *Self-duality for multi-state probabilistic cellular automata with finite range interactions*, 2002 *J. of Statistical Physics* **106** 923-930.

[13] Cox J. T. and Durrett R., *Nonlinear voter model*, 1991 In: *R. Durrett and H. Kesten (eds.), Random Walks, "Brownian Motion and Interacting Particle Systems"* Birkhäuser 189-201.

[14] Liggett T. M., *Coexistence in threshold voter models*, 1994 *Ann. Prob.* **22** 764-802.

[15] Domany E. and Kinzel W., *Equivalence of cellular automata to Ising models and direct percolation*, 1984 *Phys. Rev. Lett.* **53** 311-314.

[16] Atman A. P. F. and Moreira J. G., *Growth exponent in the Domany-Kinzel cellular automaton*, 2000 *Eur. Phys. J. B.* **16** 501-505.

[17] Kemper A., Schadschneider A. and Zittartz J., *Transfer-matrix DMRG for stochastic models: the Domany-Kinzel cellular automaton*, 2001 *J. Phys. A: Math. Gen.* **34** 279-280.

[18] Rieger H., Schadschneider A. and Schreckenberg M., *Reentrant behaviour in the Domany-Kinzel cellular automaton*, 1994 *J. Phys. A* **27** 423-430.

[19] Athreya K. B. and Ney P. E., *Branching processes*, 1972 Springer-Verlag, New York.

[20] Williams T. and Bjerknes R., *Stochastic model for abnormal clone spread through epithelial layer*, 1972 *Nature* **236** 19-22.

[21] López F. J. and Sanz G., *Stochastic comparisons for general probabilistic cellular automata*, 2000 *Stat. Prob. Lett.* **46** 401-410.

[22] Jiménez A., Tiampo K. F. and Posadas A. M., *An Ising model for earthquake dynamics*, 2007 *Nonlinear Processes in Geophysics* **14** 5-15.

[23] Kordzakhia G. and Lalley S. P., *A two-species competition model on* $\mathbb{Z}^d$, 2005 *Stochastic Processes and their Applications* **115** 781-796.

[24] Koulis T. *Stochastic modeling of sea ice*, 2003 Presented at JSM 2003, San Francisco, USA.